An Introduction to SPSS

Dr. Krishan K. Pandey

What is in this initial introduction Class

- SPSS interface: data view and variable view
- How to enter data in SPSS
- How to import external data into SPSS
- How to clean and edit data
- How to transform variables
- How to sort and select cases
- How to get descriptive statistics

Data used for this Lecture

- We use 2009 Youth Risk Behavior Surveillance System (YRBSS, CDC) as an example.
 - YRBSS monitors priority health-risk behaviors and the prevalence of obesity and asthma among youth and young adults.
 - The target population is high school students
 - Multiple health behaviors include drinking, smoking, exercise, eating habits, etc.


SPSS interface

- Data view
 - The place to enter data
 - Columns: variables
 - Rows: records
- Variable view
 - The place to enter variables
 - List of all variables
 - Characteristics of all variables


Before the data entry

- You need a code book/scoring guide
- You give ID number for each case (NOT real identification numbers of your subjects) if you use paper survey.
- If you use online survey, you need something to identify your cases.
- You also can use Excel to do data entry.


Example of a code book




Enter data in SPSS 19.0




Enter variables



Enter variables




Enter cases




Import data from Excel

- Select File → Open → Data
- Choose Excel as file type
- Select the file you want to import
- Then click Open





Open Excel files in SPSS





Import data from CVS file

- CVS is a comma-separated values file.
- If you use Qualtrics to collect data (online survey), you will get a CVS data file.
- Select File → Open → Data
- Choose **All files** as file type
- Select the file you want to import
- Then click Open




















Clean data after import data files

- Key in values and labels for each variable
- Run frequency for each variable
- Check outputs to see if you have variables with wrong values.
- Check missing values and physical surveys if you use paper surveys, and make sure they are real missing.
- Sometimes, you need to recode string variables into numeric variables


Q1 How old are you

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	1 12 years old or younger	19	.1	.1	.1
	2 13 years old	17	.1	.1	.2
	3 14 years old	1851	11.3	11.3	11.5
	4 15 years old	4045	24.7	24.7	36.3
	5 16 years old	4234	25.8	25.9	62.2
	6 17 years old	3963	24.2	24.2	86.4
	7 18 years old or older	2215	13.5	13.6	100.0
	8	1	.0	.0	100.0
	9	1	.0	.0	100.0
	Total	16344	99.6	100.0	
Mi sing	System	66	.4		
rotal		16410	100.0		


Wrong

Variable transformation

- Recode variables
 - Select Transform → Recode into Different Variables
 - 2. Select variable that you want to transform (e.g. Q20): we want 1= Yes and 0 = No
 - 3. Click Arrow button to put your variable into the right window
 - 4. Under Output Variable: type name for new variable and label, then click Change
 - 5. Click Old and New Values



- 6. Type 1 under Old Value and 1 under New Value, click Add. Then type 2 under Old Value, and 0 under New Value, click Add.
- 7. Click Continue after finish all the changes.
- 8. Click Ok




Variable transformation

- Compute variable (useYRBSS 2009 data)
 - Example 1. Create a new variable: drug_use (During the past 30 days, any use of cigarettes, alcohol, and marijuana is defined as use, else as non-use). There are two categories for the new variable (use vs. non-use). Coding: 1= Use and 0 = Non-use
 - 1. Use Q30, Q41, and Q47 from 2009 YRBSS survey
 - 2. Non-users means those who answered **0** days/times to all three questions.
 - 3. Go to Transform → Compute Variable

- 4. Type "drug_use" under Target Variable
- 5. Type "0" under Numeric Expression. 0 means

Non-use

6. Click If button.




7. With help of that Arrow button, type

Q30= 1 & Q41 = 1 & Q47= 1


then click Continue

8. Do the same thing for Use, but the numeric expression is different:

Q30> 1 | Q41> 1 | Q47>1




9. Click OK 10. After click OK, a small window asks if you want to change existing variable because drug_use was already created when you first define non-use. 11. Click ok.




- Compute variables
 - Example 2. Create a new variable drug_N that assesses total number of drugs that adolescents used during the last 30 days.
 - 1. Use Q30 (cigarettes), 41 (alcohol), 47 (marijuana), and 50 (cocaine). The number of drugs used should be between 0 and 4.
 - 2. First, recode all four variables into two categories: 0 = non-use (0 days), 1 = use (at least 1 day/time)
 - 3. Four variables have 6 or 7 categories

- 4. Recode four variables: 1 (old) = 0 (new), 2-6/7 (old) = 1 (New).
- 5. Then select Transform Compute Variable




- 6. Type drug_N under Target Variable
- 7. Numeric Expression: SUM (Q30r,Q41r,Q47r,Q50r)
- 8. Click OK




- Compute variables
 - Example 3: Convert string variable into numeric variable

- 1. Enter 1 at Numeric Expression.
- 2. Click If button and type Q2 = 'Female'
- 3. Then click Ok.
- 4. Enter 2 at Numeric Expression.
- 5. Click If button and type Q2 = 'Male'
- 6. Then click Ok




Sort and select cases

- Sort cases by variables: Data Sort Cases
- You can use Sort Cases to find missing.



Sort and select cases

- Select cases
 - Example 1. Select Females for analysis.
 - 1. Go to Data Select Cases
 - 2. Under Select: check the second one
 - 3. Click If button




4. Q2 (gender) = 1,

1 means Female


- 5. Click Continue
- 6. Click Ok

Unselected cases:
Q2 = 2




Sort and select cases

7. You will see a new variable: filter_\$ (Variable view)



Sort and select cases


- Select cases
 - Example 2. Select cases who used any of cigarettes, alcohol, and marijuana during the last 30 days.
 - 1. Data Select Cases
 - 2. Click If button
 - 3. Type Q30 $> 1 \mid Q41 > 1 \mid Q47 > 1$, click Continue




Basic statistical analysis


- Descriptive statistics
 - Purposes:
 - 1. Find wrong entries
 - 2. Have basic knowledge about the sample and targeted variables in a study
 - 3. Summarize data


Analyze → Descriptive statistics → Frequency





Frequency table





THANK YOU